

Fig. 1




Fig. 2




Fig. 3




Fig. 4




Fig. 5




Fig. 6





Fig. 7

BEST AVAILABLE COPY



— 100nm

Fig. 8A




Fig. 8B




Fig. 8C

BEST AVAILABLE COPY




Fig. 9A




Fig. 9B




Fig. 9C




Fig. 10A-1




Fig. 10A-2




Fig. 10A-3

Effects of protamine sulfate on gene transfer by HVJ envelope vector




Fig. 10B-1




Fig. 10B-2




Fig. 10B-3




Fig. 11A-1




Fig. 11A-2




Fig. 11B




Fig. 12




Fig. 13A




Fig. 13B




Fig. 14




Fig. 15




Fig. 16A




Fig. 16B




Fig. 16C

TEST AVAILABLE COPY




Fig. 16D-1





Fig. 16D-2










Fig. 16D-3



VEGF 200ng



pcNK5 6.7 μ g




pcNK5 13.3 μ g

Fig. 16E



Fig. 16F


Fluorescence image



60 min

Fig. 17A-1


Phase-contrast image



60 min

Fig. 17A-2


Fluorescence image



10 min

Fig. 17B-1


Phase-contrast image



10 min

Fig. 17B-2

BEST AVAILABLE COPY



Protamine sulfate concentration and centrifugation

Fig. 18A




Fig. 18B

(15 K rpm, 10 min, 20°C)




Fig. 18C




Fig. 19




Fig. 20




Fig. 21